



Pyridoxine Tablets BP 50mg	
<p>Composition: Each uncoated tablet contains: Pyridoxine Hydrochloride BP 50 mg. Excipients q.s.</p> <p>Indications: It is used to treat low levels of Vitamin B6 in the body.</p> <p>Properties: Pyridoxine is the 4-methanol form of vitamin B6 and is converted to pyridoxal 5-phosphate in the body. Pyridoxal 5-phosphate is a coenzyme for synthesis of amino acids, neurotransmitters (serotonin, norepinephrine), sphingolipids, aminolevulinic acid.</p> <p>Dosage / Mode of Administration: Oral Consult a physician before use. Adults: 50-150 mg daily in divided doses. Children: Not recommended for children</p> <p>Side Effects: The effects are headache, nausea, drowsiness and allergic reactions.</p> <p>Interactions: It reduces the effects of levo dopa. Some products that may interact with this vitamin include Altretamine, Cisplatin, Phenytoin.</p> <p>Precautions: Consult doctor if any of these medications is co-administered; levodopa, isoniazid, oral contraceptives and penicillamine.</p> <p>Storage: Store in cool dry place. Keep below 25°C. Protect from direct sunlight and humidity. Keep out of the reach of children.</p> <p>Presentation: 1x10, 2x10, 3x10, 10x10 blisters of 10 tablets in 1 carton and 100x10-100 blisters of 10 tablets in 1 carton, For HDPE containers 100/500/1000 tablets in 1 jar.</p>	
 Parcela 726, Avenida das Industrias, Machava Maputo-Mozambique	

112 x 112mm

Piridoxina Comprimidos BP 50mg	
<p>Composição: Cada comprimido não revestido contém: Cloridrato de Piridoxina BP 50 mg. Excipientes q.s.</p> <p>Indicações: É usado no tratamento de níveis baixos de Vitamina B6 no corpo.</p> <p>Propriedades: Piridoxina é a forma metanol-4 da vitamina B6 e é convertida no corpo para fosfato piridoxal-5. O fosfato piridoxal-5 é uma coenzima para a síntese dos aminoácidos, neurotransmissores (serotonina, norepinefrina), esfingolípídeos, ácido aminolevulínico.</p> <p>Dosagem / Modo de Administração: Oral Consulte um médico antes de usar. Adultos: 50-150 mg por dia em doses divididas. Crianças: Não é recomendado para crianças</p> <p>Efeitos Colaterais: Os efeitos secundários são dores de cabeça, náuseas, sonolência e reações alérgicas.</p> <p>Interações medicamentosas: Reduz os efeitos de levodopa. Alguns produtos que podem interagir com esta vitamina incluem Altretamina, Cisplatina e Fenitoína.</p> <p>Precauções: Consulte o seu médico se for coadministrado algum destes medicamentos; levodopa, isoniazida, contraceptivos orais e penicilamina.</p> <p>Conservação: Armazenar em local fresco e seco. Mantenha abaixo de 25°C. Proteger da luz solar direta e da humidade. Manter fora do alcance das crianças.</p> <p>Apresentação: 1x10, 2x10, 3x10, 10x10 10 blisters de 10 comprimidos em 1 caixa e 100x10-100 blisters de 10 comprimidos em caixa. Frasco de HDPE com 100/500/1000 comprimidos em 1 frasco.</p>	
 Parcela 726, Avenida das Industrias, Machava Maputo-Mozambique	

112 x 112mm

ARTWORK DETAIL LABEL

Product	Pyridoxine Tablets BP 50 mg		
Buyer / Country	Mozambique	Component	Leaflet
Dimension	112 x 112mm	Pack	NA
New Item Code	STR/M/PI/033	Old Item code	NA
Colour Shades	██████ BLACK	No. of Colours	1

Change Control No.			Artwork Version	R0
Design/Style	Front & Back Printing			
Substrate	60 gsm maplitho paper			
Special Instruction	Colours to be matched as per pantone shades only. Printing clarity to be clear and sharp			
Autocartonater Requirement	NA			
Prepared By:	Reviewed By:	Reviewed By:	Reviewed By:	Approved By:
PDC-AW	Mkt/BD	RAD	PD	QA

FOR ARTWORK REVISION PURPOSE

To be filled by SCM (Planning)	Effective Date :		Sign/ Date:	
--------------------------------	------------------	--	-------------	--

Caution to the printer: Before processing please ensure that the ARTWORK received for printing is exactly in line with APPROVED ARTWORK provided to you. In case of any FONTS/DESIGN are Mis - matching with the APPROVED ARTWORK, please inform to PDC for further action. **DO NOT MAKE ANY CHANGE TO THE ARTWORK WITHOUT WRITTEN INSTRUCTION FROM PDC.**

QA/068/F-01/R0